

Combinatorial Bethe ansatz and ultradiscrete Riemann theta function with rational characteristics ¹

Atsuo Kuniba and Reiho Sakamoto

ABSTRACT. The $U_q(\widehat{sl}_2)$ vertex model at $q = 0$ with periodic boundary condition is an integrable cellular automaton in one-dimension. By the combinatorial Bethe ansatz, the initial value problem is solved for arbitrary states in terms of an ultradiscrete analogue of the Riemann theta function with rational characteristics.

1. INTRODUCTION

Solvable vertex models [2] associated with a quantum affine algebra U_q yield completely integrable one-dimensional cellular automata at $q = 0$ [9, 8, 6]. The simplest case of $U_q(\widehat{sl}_2)$ with periodic boundary condition is known as the periodic box-ball system [17, 22]. Here is an example of the time evolution pattern:

```

t = 0:  11122222111122222222111111111111112222111211
t = 1:  11111111222211111111222222221111111111222122
t = 2:  2222111111112222111111111111222222221111211
t = 3:  111122222222111122221111111111111111112222122
t = 4:  22221111111122221111222222221111111111111211
t = 5:  11112222111111112222111111112222222211111121
t = 6:  22221111222211111111222211111111111111222212
t = 7:  111122221111222222221111222221111111111111121
t = 8:  11111111222211111111222211112222222211111112
t = 9:  22221111111122221111111122221111111111222221

```

The dynamics is described as a motion of balls (letter 2) hopping to the right along the periodical array of boxes. The fusion transfer matrices $T_1, T_2, \dots$ in the $q = 0$ vertex model give rise to a commuting family of deterministic time evolutions. The above example is the evolution under T_9 .

In [17], the initial value problem of the periodic box-ball system is solved by an inverse scattering method. It synthesizes the combinatorial versions of the Bethe ansatz [3] at $q = 1$ [13] and $q = 0$ [14]. The action-angle variables are constructed from the rigged configurations ($q = 1$) by imposing a certain equivalence relations specified by the string center equation ($q = 0$). The time evolution $T_t^t(p)$ of any state p is obtained by an explicit algorithm whose computational steps are independent of t .

The Bethe ansatz approach [17] captures several characteristic features in the quasi-periodic solutions of soliton equations [4, 5, 11]. It generalizes the connection between the original box-ball system on infinite lattice [20] and soliton solutions via the ultradiscretization [21] to a periodic setting. For instance, the nonlinear dynamics of the periodic box-ball system becomes a straight motion of the Bethe roots (angle variable) which live in an ultradiscrete analogue (4.1) of the Jacobi variety.

¹ Mathematics Subject Classification (2000). 81R50, 82B23, 37B15, 37K15, 05E15
Keywords: crystal basis, periodic box-ball system, Bethe ansatz, Riemann theta function

In this paper we express the algorithmic solution of the initial value problem [17] by an explicit formula involving the ultradiscretization of the Riemann theta function with rational characteristics ($\mathbf{z} \in \mathbb{R}^g$, $\mathbf{a} \in (\mathbb{Q}/\mathbb{Z})^g$):

$$(1.1) \quad \begin{aligned} \Theta_{\mathbf{a}}(\mathbf{z}) &= \lim_{\epsilon \rightarrow +0} \epsilon \log \vartheta_{\mathbf{a}}(\mathbf{z}) \\ &= - \min_{\mathbf{n} \in \mathbb{Z}^g} \{ {}^t(\mathbf{n} + \mathbf{a})\Omega(\mathbf{n} + \mathbf{a})/2 + {}^t(\mathbf{n} + \mathbf{a})\mathbf{z} \}. \end{aligned}$$

Here $\vartheta_{\mathbf{a}}(\mathbf{z})$ is the Riemann theta function [19] with a pure imaginary period matrix:

$$(1.2) \quad \vartheta_{\mathbf{a}}(\mathbf{z}) = \sum_{\mathbf{n} \in \mathbb{Z}^g} \exp\left(-\frac{{}^t(\mathbf{n} + \mathbf{a})\Omega(\mathbf{n} + \mathbf{a})/2 + {}^t(\mathbf{n} + \mathbf{a})\mathbf{z}}{\epsilon}\right).$$

Here Ω is a symmetric positive definite integer matrix (5.1). The function $\Theta_{\mathbf{a}}(\mathbf{z})$ enjoys the quasi-periodicity analogous to the Riemann theta function:

$$(1.3) \quad \Theta_{\mathbf{a}}(\mathbf{z} + \mathbf{v}) = \Theta_{\mathbf{a}}(\mathbf{z}) + {}^t\mathbf{v}\Omega^{-1}(\mathbf{z} + \mathbf{v}/2) \quad \text{for any } \mathbf{v} \in \Omega\mathbb{Z}^g.$$

Our main Theorem 5.1 covers *all* the states in the periodic box-ball system, which generalizes the earlier result [15] on those states in which the amplitudes of solitons were assumed to be distinct. The theorem is derived from the recently obtained piecewise linear formula [16] for the Kerov-Kirillov-Reshetikhin (KKR) bijection [13] from rigged configurations to highest paths. It involves the ultradiscretization of the tau function for the KP hierarchy [10]. A state p of the periodic box-ball system is effectively treated as the infinite system $p \otimes p \otimes p \otimes \dots$, and the ultradiscrete tau function for such states turns out to be expressible in terms of $\Theta_{\mathbf{a}}(\mathbf{z})$ as in (5.5).

In section 2 we define the periodic box-ball system. In section 3 various data in the combinatorial Bethe ansatz at $q = 1$ [13] and $q = 0$ [14] are introduced. In section 4 we recall the solution of the initial value problem by the inverse scattering transform. In section 5 we present our main Theorem 5.1 in this paper.

2. PERIODIC BOX-BALL SYSTEM

Let us recall the periodic box-ball system without getting much into the crystal base theory [12]. For a positive integer l , let $B_l = \{(x_1, x_2) \in (\mathbb{Z}_{\geq 0})^2 \mid x_1 + x_2 = l\}$ and set $u_l = (l, 0) \in B_l$. The element (x_1, x_2) will also be denoted by the sequence

$\overbrace{1 \dots 1}^{x_1} \overbrace{2 \dots 2}^{x_2}$, which is the semistandard Young tableau (without a frame) containing

the letter i x_i times. In this notation $u_l = \overbrace{1 \dots 1}^l$, and the two elements $(1, 0)$ and $(0, 1)$ in B_1 are simply written as 1 and 2, respectively. In the following, the symbol $\otimes$ meaning the tensor product of crystals can just be understood as a product of sets. Define the map $R : B_l \otimes B_1 \rightarrow B_1 \otimes B_l$ by

$$\begin{aligned} (x_1, x_2) \otimes 1 &\mapsto \begin{cases} 1 \otimes (l, 0) & \text{if } (x_1, x_2) = (l, 0) \\ 2 \otimes (x_1 + 1, x_2 - 1) & \text{otherwise,} \end{cases} \\ (x_1, x_2) \otimes 2 &\mapsto \begin{cases} 2 \otimes (0, l) & \text{if } (x_1, x_2) = (0, l) \\ 1 \otimes (x_1 - 1, x_2 + 1) & \text{otherwise.} \end{cases} \end{aligned}$$

R is a bijection and called the combinatorial R . We write the relation $R(u \otimes b) = b' \otimes u'$ simply as $u \otimes b \simeq b' \otimes u'$, and similarly for any consequent relation of the form $a \otimes u \otimes b \otimes c \simeq a \otimes b' \otimes u' \otimes c$.

A state of the periodic box-ball system is an array of 1 and 2, which is regarded as an element $b_1 \otimes \cdots \otimes b_L \in B_1^{\otimes L}$ with L being the system size. Let the number of $2 \in B_1$ appearing in $b_1 \otimes \cdots \otimes b_L$ be N . We assume $L \geq 2N$. The other case can be reduced to this case by the exchange $1 \leftrightarrow 2$ as explained in §3.3 of [17]. Let $\mathcal{P}$ be the set of such states. Then the time evolution $T_l : \mathcal{P} \rightarrow \mathcal{P}$ is defined by

$$(2.1) \quad u_l \otimes p \simeq p^* \otimes v_l, \quad v_l \otimes p \simeq T_l(p) \otimes v_l.$$

In the first relation, one applies the combinatorial R L times to carry u_l through $p \in \mathcal{P}$ to the right. It determines $v_l \in B_l$ and $p^* \in \mathcal{P}$ uniquely. (p^* will play no role in the sequel.) Then the second relation using the so obtained v_l specifies $T_l(p)$, where the appearance of the same v_l in the right hand side is a non-trivial claim ([17] (2.10)). v_l is dependent on p as opposed to u_l .

The combinatorial R is the identity map on $B_1 \otimes B_1$, therefore T_1 is just the cyclic shift $T_1(b_1 \otimes \cdots \otimes b_L) = b_L \otimes b_1 \otimes \cdots \otimes b_{L-1}$. The commutativity $T_l T_k = T_k T_l$ holds for any k, l ([17] Theorem 2.2).

Example 2.1. We depict the relation $R(u \otimes b) = b' \otimes u'$ by a vertex diagram:

$$\begin{array}{c} b \\ | \\ u \text{---} \text{---} u' \\ | \\ b' \end{array}$$

Then for $p = 1121221 \in B_1^{\otimes 7}$ (symbol $\otimes$ omitted), one has $v_2 = (1, 1) = 12 \in B_2$ from (2.1). The time evolution $T_2(1121221)$ is obtained by the composition of the vertex diagrams

$$\begin{array}{ccccccc} 1 & 1 & 2 & 1 & 2 & 2 & 1 \\ 12 \text{---} \text{---} 11 \text{---} \text{---} 11 \text{---} \text{---} 12 \text{---} \text{---} 11 \text{---} \text{---} 12 \text{---} \text{---} 22 \text{---} \text{---} 12 \\ 2 & 1 & 1 & 2 & 1 & 1 & 2 \end{array}$$

leading to $T_2(1121221) = 2112112$.

3. COMBINATORIAL BETHE ANSATZ DATA

Let $\mathcal{P}_+$ be the subset of $\mathcal{P}$ consisting of the sl_2 -highest states. Namely, $\mathcal{P}_+ = \{p \in \mathcal{P} \mid \tilde{e}_1 p = 0\}$, where $\tilde{e}_1$ is a Kashiwara operator [12]. In practice, $p = b_1 \otimes \cdots \otimes b_L$ is highest if and only if $b_1 \otimes \cdots \otimes b_k$ has nonnegative weights for any $1 \leq k \leq L$.

According to the combinatorial Bethe ansatz at $q = 1$ [13], there is one to one correspondence between $\mathcal{P}_+$ and the set of the objects called the *rigged configurations*. We let $\phi(p_+) = (\mu, J)$ denote the Kerov-Kirillov-Reshetikhin (KKR) bijection. Here p_+ is highest state and the right hand side is a rigged configuration. The data μ , called a configurations, is just a Young diagram for a partition of N , for which $N \leq L/2$ is assumed.

In what follows, we let $\mathbb{I} = \{i_1 < i_2 < \cdots < i_g\}$ be the set of distinct lengths of rows of μ , and the multiplicity of the length $i (\in \mathbb{I})$ rows by $m_i (\geq 1)$. In terms of the non-increasing array of parts we have

$$(3.1) \quad \mu = (i_g^{m_{i_g}} \dots i_1^{m_{i_1}}).$$

Obviously, a configuration μ is equivalent with the data $\mathbb{I}$ and $m_{i_1}, \dots, m_{i_g}$.

To each row of μ , there is assigned a nonnegative integer called a rigging, which is represented by the symbol J in (μ, J) . See Example 5.2. Let $J_{i,1}, \dots, J_{i,m_i}$ be the ones assigned to the m_i rows of length i in μ from the bottom to the top. The data

(μ, J) with $J = \{(J_{i,\alpha}) \mid i \in \mathbb{I}, 1 \leq \alpha \leq m_i\}$ will be called a rigged configuration if $0 \leq J_{i,1} \leq \dots \leq J_{i,m_i} \leq p_i$ holds for each $i \in \mathbb{I}$. Here p_i is defined by

$$(3.2) \quad p_i = L - 2 \sum_{j \in \mathbb{I}} \min(i, j) m_j,$$

which is nonnegative if μ is a configuration.

We introduce the index set

$$(3.3) \quad \tilde{\mathbb{I}} = \{(i, \alpha) \mid i \in \mathbb{I}, 1 \leq \alpha \leq m_i\}, \quad \gamma = |\tilde{\mathbb{I}}| = m_{i_1} + \dots + m_{i_g}$$

labeling the rows of μ . The following vectors and matrix whose components are labeled with $\tilde{\mathbb{I}}$ will be used:

$$(3.4) \quad \tilde{\mathbf{h}}_j = (\min(i, j))_{(i, \alpha) \in \tilde{\mathbb{I}}} \in \mathbb{Z}^\gamma \quad (j \in \mathbb{I}),$$

$$(3.5) \quad \tilde{\mathbf{J}} = (J_{i,\alpha} + \alpha - 1)_{(i, \alpha) \in \tilde{\mathbb{I}}} \in \mathbb{Z}^\gamma,$$

$$(3.6) \quad A = (A_{i\alpha, j\beta})_{(i, \alpha), (j, \beta) \in \tilde{\mathbb{I}}}, \quad A_{i\alpha, j\beta} = \delta_{i,j} \delta_{\alpha, \beta} (p_i + m_i) + 2 \min(i, j) - \delta_{i,j}.$$

We also introduce the vectors and matrices whose indices range over the previously introduced set $\mathbb{I} = \{i_1, \dots, i_g\}$ as

$$(3.7) \quad \mathbf{h}_j = (\min(i, j))_{i \in \mathbb{I}} \in \mathbb{Z}^g,$$

$$(3.8) \quad \mathbf{p} = (p_i)_{i \in \mathbb{I}} = L \mathbf{h}_1 - 2 \sum_{j \in \mathbb{I}} m_j \mathbf{h}_j \in \mathbb{Z}^g,$$

$$(3.9) \quad \mathbf{J} = (J_{i,1} + \dots + J_{i,m_i})_{i \in \mathbb{I}} \in \mathbb{Z}^g,$$

$$(3.10) \quad F = (F_{i,j})_{i,j \in \mathbb{I}}, \quad F_{i,j} = \sum_{\beta=1}^{m_j} A_{i\alpha, j\beta} = \delta_{ij} p_i + 2 \min(i, j) m_j,$$

$$(3.11) \quad M = \text{diag}(m_i)_{i \in \mathbb{I}},$$

The matrices A, F were introduced in the study of $q = 0$ Bethe equation [14]. In particular, A is the coefficient matrix of the linearized Bethe equation (string center equation), which is known to be positive definite. The result of the sum in (3.10) is independent of α .

4. ACTION-ANGLE VARIABLE AND INVERSE SCATTERING TRANSFORM

Any state $p \in \mathcal{P}$ can be made highest by applying the cyclic shift appropriately. Namely, one can express p as $p = T_1^d(p_+)$ with some integer d and highest state $p_+ \in \mathcal{P}_+$. Although the choice of d and p_+ is not unique, the configuration of $\phi(p_+)$ is unique. The set of states is decomposed according to the configuration of p_+ as $\mathcal{P} = \sqcup_\mu \mathcal{P}(\mu)$. We call μ the action variable. It is a conserved quantity ([17] Lemma C.3, Corollary 3.5), therefore each subset $\mathcal{P}(\mu)$ is invariant under any time evolution T_l . The action variable μ is equivalent with the data $\mathbb{I} = \{i_1, \dots, i_g\}$ and $\{m_i \mid i \in \mathbb{I}\}$ via (3.1). In the periodic box-ball system, $\mathbb{I}$ is the list of amplitudes of solitons and m_i is the number of solitons with amplitude i .

Given an action variable μ , the set of angle variable is given by

$$(4.1) \quad \mathcal{J}(\mu) = (\mathcal{I}_{m_{i_1}} \times \dots \times \mathcal{I}_{m_{i_g}}) / \Gamma, \quad \Gamma = A \mathbb{Z}^\gamma,$$

where $\mathcal{I}_n = (\mathbb{Z}^n - \Delta_n) / \mathfrak{S}_n$ is the n -dimensional lattice without the ‘‘diagonal points’’ $\Delta_n = \{(z_1, \dots, z_n) \in \mathbb{Z}^n \mid z_\alpha = z_\beta \text{ for some } 1 \leq \alpha \neq \beta \leq n\}$ divided by the action of the symmetric group $\mathfrak{S}_n$. Plainly, it is the set of array of n distinct

integers whose ordering does not matter. In the sequel, elements in $\mathcal{I}_{m_{i_1}} \times \cdots \times \mathcal{I}_{m_{i_g}}$ and its image in $\mathcal{J}(\mu)$ will be denoted by the same symbol. The set $\mathcal{J}(\mu)$ is finite and its cardinality is given by [14, 17]

$$(4.2) \quad |\mathcal{J}(\mu)| = (\det F) \prod_{i \in \mathbb{I}} \frac{1}{m_i} \binom{p_i + m_i - 1}{m_i - 1} = \frac{L}{p_{i_g}} \prod_{i \in \mathbb{I}} \binom{p_i + m_i - 1}{m_i}.$$

This is a positive integer if μ is a configuration. (When $|\mu| = L/2$, we have $p_{i_g} = 0$. In this case, the factor $\frac{L}{p_{i_g}} \binom{p_{i_g} + m_{i_g} - 1}{m_{i_g}}$ is to be understood as $\frac{L}{m_{i_g}}$.) The formula (4.2) tells the number of states containing m_i solitons with amplitude i . The completeness of soliton states is known in the sense that $\sum_{\mu} |\mathcal{J}(\mu)| = \binom{L}{N}$ for $0 \leq N \leq L/2$, where the sum extends over those m_i such that $\sum_i i m_i = N$ ([14], Theorems 3.5 and 4.9). The states in section 1 have the length $L = 45$ and μ depicted in Example 5.2. For this μ and L , the formula (4.2) tells $|\mathcal{J}(\mu)| = 316350$.

Let us specify the angle variable for a given state $p \in \mathcal{P}(\mu)$. Suppose p is expressed as $p = T_1^d(p_+)$ in terms of a highest state p_+ and the cyclic shift T_1 . Let the image of the KKR bijection be $\phi(p_+) = (\mu, J)$. From the configuration μ we know $\mathbb{I} = \{i_1 < i_2 < \cdots < i_g\}$ and $m_{i_1}, \dots, m_{i_g}$ by (3.1). Then the angle variable is found by applying the definitions (3.4), (3.5), (3.6) to $J = (J_{i,\alpha})_{(i,\alpha) \in \mathbb{I}}$ as

$$(4.3) \quad \begin{aligned} \Phi: \mathcal{P}(\mu) &\longrightarrow \mathbb{Z} \times \mathcal{P}_+ \longrightarrow \mathcal{J}(\mu) \\ p &\longmapsto (d, p_+) \longmapsto (\tilde{\mathbf{J}} + d\tilde{\mathbf{h}}_1)/\Gamma. \end{aligned}$$

Due to the shift $+\alpha - 1$ in (3.5) and the rigging condition $0 \leq J_{i,1} \leq \cdots \leq J_{i,m_i} \leq p_i$, it follows that the components of $\tilde{\mathbf{J}}$ hence $\tilde{\mathbf{J}} + d\tilde{\mathbf{h}}_1$ within each block labeled by $i \in \mathbb{I}$ are distinct. In (4.3) we regard the $\tilde{\mathbf{J}} + d\tilde{\mathbf{h}}_1$ as a representative of an element in $\mathcal{J}(\mu)$. The non-uniqueness of d, p_+ is cancelled by mod Γ making Φ well-defined ([17] Proposition 3.7).

Theorem 4.1 ([17] Theorem 3.12). *Φ is a bijection and the following diagram is commutative.*

$$\begin{array}{ccc} \mathcal{P}(\mu) & \xrightarrow{\Phi} & \mathcal{J}(\mu) \\ T_l \downarrow & & \downarrow T_l \\ \mathcal{P}(\mu) & \xrightarrow{\Phi} & \mathcal{J}(\mu) \end{array}$$

Here the time evolution on $\mathcal{J}(\mu)$ is defined by $T_l(\tilde{\mathbf{J}}) = \tilde{\mathbf{J}} + \tilde{\mathbf{h}}_l$.

The composition $\Phi^{-1} \circ T_l \circ \Phi$ achieves the solution of the initial value problem by the inverse scattering transform. Namely, $\Phi^{\pm 1}$ is the direct/inverse scattering map which transforms the dynamics T_l of the periodic box-ball system into the straight motion on $\mathcal{J}(\mu)$ with the velocity $\tilde{\mathbf{h}}_l$. In this sense, $\mathcal{J}(\mu)$ serves as an ultradiscrete analogue of the Jacobi variety in quasi-periodic solutions to soliton equations [4, 5, 11].

By using Theorem 4.1, a closed formula for the fundamental period, i.e., the smallest positive integer such that $T_l^N(p) = p$ for any l and p has been obtained in Theorem 4.9 of [17], which supplements a recursive description of the $l = \infty$ case [22]. The states in section 1 have the fundamental period 3515. See also [18] for a generalization to $A_n^{(1)}$ based on the Bethe eigenvalue at $q = 0$.

We note that the time evolution of the angle variable $\tilde{\mathbf{J}}$ (3.5) leads to the rule

$$(4.4) \quad T_l(\mathbf{J}) = \mathbf{J} + M\mathbf{h}_l$$

for the “bundled” angle variable $\mathbf{J}$ (3.9), where M is defined by (3.11).

5. EXPLICIT FORMULA FOR THE SOLUTION OF THE INITIAL VALUE PROBLEM

Fix the action variable μ (3.1) and set

$$(5.1) \quad \Omega = MF = (\Omega_{i,j})_{i,j \in \mathbb{I}}, \quad \Omega_{i,j} = \delta_{ij}p_i m_i + 2 \min(i, j) m_i m_j,$$

where F is defined in (3.11). Ω is a positive definite symmetric integer matrix which will play the role of the period matrix. Note that

$$(5.2) \quad \Omega \mathbf{h}_1 = LM\mathbf{h}_1.$$

In addition to the ultradiscrete Riemann theta function $\Theta_{\mathbf{a}}(\mathbf{z})$ in (1.1), we need $\chi(\mathbf{s}; \tilde{\mathbf{I}})$. It is a function of $\mathbf{s} = (s_i)_{i \in \mathbb{I}} \in \mathbb{Z}^g$ as well as $\tilde{\mathbf{I}} = (I_{i,\alpha} + \alpha - 1)_{(i,\alpha) \in \tilde{\mathbb{I}}} \in \mathcal{I}_{m_{i_1}} \times \cdots \times \mathcal{I}_{m_{i_g}}$ that takes rational values:

$$(5.3) \quad \chi(\mathbf{s}; \tilde{\mathbf{I}}) = \chi(\mathbf{s}'; \tilde{\mathbf{I}}) \text{ if } \mathbf{s} \equiv \mathbf{s}' \pmod{M\mathbb{Z}^g},$$

$$(5.4) \quad \chi(\mathbf{s}; \tilde{\mathbf{I}}) = \sum_{i \in \mathbb{I}} \frac{1}{m_i} \sum_{1 \leq \alpha \leq s_i < \beta \leq m_i} (I_{i,\beta} - I_{i,\alpha} - \frac{p_i}{2})$$

if $0 \leq s_i < m_i$, $I_{i,1} \leq \cdots \leq I_{i,m_i}$ for all $i \in \mathbb{I}$.

Here the second sum in (5.4) extends over the pairs (α, β) . In particular, $\chi(\mathbf{0}; \tilde{\mathbf{I}}) = 0$. The condition $I_{i,1} \leq \cdots \leq I_{i,m_i}$ is needed to cope with the $\mathfrak{S}_{m_i}$ symmetry in $\mathcal{I}_{m_i}$.

Given $\mathbf{I} \in \mathbb{Z}^g$ and $\tilde{\mathbf{I}} = (I_{i,\alpha} + \alpha - 1)_{(i,\alpha) \in \tilde{\mathbb{I}}} \in \mathcal{I}_{m_{i_1}} \times \cdots \times \mathcal{I}_{m_{i_g}}$, we define the ultradiscrete tau function by

$$(5.5) \quad \tau_r(k) = \max_{\mathbf{s} \in \mathbb{Z}^g / M\mathbb{Z}^g} \left\{ \Theta_{M^{-1}\mathbf{s}} \left(\mathbf{I} + M(r\mathbf{h}_\infty - k\mathbf{h}_1 - \frac{\mathbf{p}}{2}) \right) + \chi(\mathbf{s}; \tilde{\mathbf{I}}) \right\}$$

for $r \in \{0, 1\}$, $k \in \mathbb{Z}$. Here $\mathbf{h}_j$, $\mathbf{p}$ and M are those in (3.7), (3.8) and (3.11).

Let us specify $\tilde{\mathbf{I}}$ and $\mathbf{I}$ in (5.5) from a given state $p \in \mathcal{P}(\mu)$. As for $\tilde{\mathbf{I}}$, it is taken to be the angle variable of $p \in \mathcal{P}(\mu)$ according to (4.3), and $\mathbf{I}$ is determined from it as follows:

$$(5.6) \quad \tilde{\mathbf{I}} = \Phi(p) = (I_{i,\alpha} + \alpha - 1)_{(i,\alpha) \in \tilde{\mathbb{I}}} \in \mathcal{J}(\mu),$$

$$(5.7) \quad \mathbf{I} = (I_{i,1} + \cdots + I_{i,m_i})_{i \in \mathbb{I}} \in \mathbb{Z}^g / F\mathbb{Z}^g.$$

This is the same relation as (3.5) and (3.9). Note that the elements $\tilde{\mathbf{I}}$ and $\mathbf{I}$ are defined mod $\Gamma = A\mathbb{Z}^g$ and mod $F\mathbb{Z}^g$, respectively, which is consistent with the relation (3.10) between A and F . Now we state the main result of this paper.

Theorem 5.1. *The state $p \in \mathcal{P}(\mu)$ corresponding to the action variable μ and the angle variable $\tilde{\mathbf{I}}$ is expressed as $p = (1 - x_1, x_1) \otimes \cdots \otimes (1 - x_L, x_L)$, where $x_k = 0, 1$ is given by*

$$(5.8) \quad x_k = \tau_0(k) - \tau_0(k-1) - \tau_1(k) + \tau_1(k-1).$$

The proof is similar to section 3 in [15]. Combined with Theorem 4.1, Theorem 5.1 leads to an explicit formula for the solution of the initial value problem. The state $(\prod_l T_l^{c_l})(p)$ is obtained just by replacing the angle variable as

$$(5.9) \quad \tilde{\mathbf{I}} \mapsto \tilde{\mathbf{I}} + \sum_l c_l \tilde{\mathbf{h}}_l$$

in (5.5). Under the change (5.9), the function $\chi(\mathbf{s}; \tilde{\mathbf{I}})$ (5.4) is invariant since it only depends on the difference $I_{i,\beta} - I_{i,\alpha}$. On the other hand, the variable $\mathbf{I}$ entering (5.5) changes as

$$(5.10) \quad \mathbf{I} \mapsto \mathbf{I} + M \left(\sum_l c_l \mathbf{h}_l \right)$$

according to the rule (4.4). Especially, $T_1^L(\mathbf{I}) = \mathbf{I} + LM\mathbf{h}_1 = \mathbf{I} + \Omega\mathbf{h}_1$ by (5.2). Thus the tau function only exhibits the change due to the quasi-periodicity (1.3) leaving (5.8) invariant. This is consistent with the fact that the cyclic shift T_1^L is trivial and (5.8) is a period L function of k . We remark that our tau function (5.5) is always integer-valued.

From (3.7) we see $\mathbf{h}_l = \mathbf{h}_\infty$ if $l \geq \max \mathbb{I} (= i_g)$. Therefore the effect (5.10) of the time evolution T_l is $\mathbf{I} \mapsto \mathbf{I} + M\mathbf{h}_\infty$, which is equivalent to setting $r \mapsto r + 1$ in (5.5). In other words, r serves as the time variable for such T_l , i.e., $T_l^t(p) = (1 - x_{1,t}, x_{1,t}) \otimes \cdots \otimes (1 - x_{L,t}, x_{L,t})$ with

$$(5.11) \quad x_{k,t} = \tau_t(k) - \tau_t(k-1) - \tau_{t+1}(k) + \tau_{t+1}(k-1)$$

holds for $l \geq \max \mathbb{I}$.

Example 5.2. Let us consider the $t = 0, 1$ states in section 1:

$$(5.12) \quad \begin{aligned} p &= 111222221111222222221111111111112222111211 \\ T_9(p) &= 11111112222111111122222222111111111222122, \end{aligned}$$

which are elements in $B_1^{\otimes 45}$. We have $p = T_1^{29}(p_+)$ with a highest state

$$p_+ = 1111122221112111122222111122222222111111111$$

which corresponds to the rigged configuration:

$$\begin{array}{|c|c|c|c|c|c|c|c|} \hline & & & & & & & \\ \hline & & & & & & & \\ \hline & & & & & & & \\ \hline & & & & & & & \\ \hline & & & & & & & \\ \hline & & & & & & & \\ \hline \end{array} \begin{array}{l} 0 \\ 6 \\ 2 \\ 10 \end{array}$$

Thus $\mathbb{I} = \{1, 4, 9\}$, $M = \text{diag}(m_1, m_4, m_9) = \text{diag}(1, 2, 1)$. $I_{i,\alpha}$ in (5.6) reads $(I_{1,1}, I_{4,1}, I_{4,2}, I_{9,1}) = (39, 31, 35, 29)$. The data $\mathbf{p}$ (3.8), $\mathbf{I}$ (5.7) and Ω (5.1) are given by

$$\mathbf{p} = \begin{pmatrix} p_1 \\ p_4 \\ p_9 \end{pmatrix} = \begin{pmatrix} 37 \\ 19 \\ 9 \end{pmatrix}, \quad \mathbf{I} = \begin{pmatrix} 39 \\ 66 \\ 29 \end{pmatrix}, \quad \Omega = \begin{pmatrix} 39 & 4 & 2 \\ 4 & 70 & 16 \\ 2 & 16 & 27 \end{pmatrix}.$$


The argument of the tau function (5.5) reads

$$(5.13) \quad \mathbf{I} + M(r\mathbf{h}_\infty - k\mathbf{h}_1 - \frac{\mathbf{p}}{2}) = \begin{pmatrix} r - k + 41/2 \\ 8r - 2k + 47 \\ 9r - k + 49/2 \end{pmatrix} =: \mathbf{z}_{r,k}.$$

The function $\chi(\mathbf{s}; \tilde{\mathbb{I}})$ in (5.4) is non-vanishing at $\mathbf{s} = (0, 1, 0)$ taking the value $\frac{1}{m_4}(I_{4,2} - I_{4,1} - \frac{p_4}{2}) = -\frac{11}{4}$. Thus the tau function is given by

$$\tau_r(k) = \max\{\Theta_{0,0,0}(\mathbf{z}_{r,k}), \Theta_{0,\frac{1}{2},0}(\mathbf{z}_{r,k}) - \frac{11}{4}\}.$$

The time evolution pattern in section 1 is reproduced according to (5.11). Here is the plot of $\tau_r(k)$ for $0 \leq r \leq 2$ and $0 \leq k \leq 45$.



The boundary values are $(\tau_0(0), \tau_1(0), \tau_2(0)) = (15, 28, 41)$ and $(\tau_0(45), \tau_1(45), \tau_2(45)) = (13, 8, 3)$. In the plot of $\tau_t(k)$ or $\tau_{t+1}(k)$, one observes a change of the gradient at those k corresponding to the front or tail of solitons in $T_9^t(p)$ (5.12).

The tau function (5.5) simplifies considerably if there are no solitons with the same amplitude, namely, $\forall m_i = 1$. We have $\chi(\mathbf{s}; \tilde{\mathbf{I}}) = 0$, $\tilde{\mathbf{I}} = \mathbf{I}$ and $\Omega = F = (\delta_{ij}p_i + 2 \min(i, j))_{i,j \in \mathbb{I}}$. Let us write the ultradiscrete Riemann theta function with characteristics $\mathbf{0}$ as $\Theta(\mathbf{z}) = \Theta_{\mathbf{0}}(\mathbf{z})$. Then $\tau_r(k) = \Theta(\mathbf{I} - \frac{\mathbf{p}}{2} - k\mathbf{h}_1 + r\mathbf{h}_{\infty})$. Consequently (5.8) reduces to

$$\begin{aligned} x_k &= \Theta(\mathbf{I} - \frac{\mathbf{p}}{2} - k\mathbf{h}_1) - \Theta(\mathbf{I} - \frac{\mathbf{p}}{2} - (k-1)\mathbf{h}_1) \\ &\quad - \Theta(\mathbf{I} - \frac{\mathbf{p}}{2} - k\mathbf{h}_1 + \mathbf{h}_{\infty}) + \Theta(\mathbf{I} - \frac{\mathbf{p}}{2} - (k-1)\mathbf{h}_1 + \mathbf{h}_{\infty}), \end{aligned}$$

which was obtained in [15]. This has the same structure as the ultradiscretization of the solution of the periodic Toda lattice by Date and Tanaka [4].

Acknowledgments The authors thank Rei Inoue for a discussion. R.S. is a research fellow of the Japan Society for the Promotion of Science. He thanks Miki Wadati for continuous encouragements.

REFERENCES

- [1] Ablowitz, M. J. and Segur, H.: Solitons and the inverse scattering transform, SIAM Studies in Appl. Math. 4. Philadelphia Pa. (1981)
- [2] Baxter, R. J.: Exactly solved models in statistical mechanics, Academic Press, London (1982)
- [3] Bethe, H. A.: Zur Theorie der Metalle, I. Eigenwerte und Eigenfunktionen der linearen Atomkette, Z. Physik **71** 205–231 (1931)
- [4] Date, E. and Tanaka, S.: Periodic multi-soliton solutions of Korteweg-de Vries equation and Toda lattice, Prog. Theoret. Phys. Suppl. **59** 107–125 (1976)

- [5] Dubrovin, B. A., Matveev, V. B. and Novikov, S. P.: Nonlinear equations of Korteweg-de Vries type, finite-band linear operators and Abelian varieties Russian Math. Surveys **31** 59–146 (1976)
- [6] Fukuda, K., Okado, M. and Yamada, Y.: Energy functions in box ball systems, Int. J. Mod. Phys. A **15** 1379–1392 (2000)
- [7] Gardner, C. S., Greene, J. M., Kruskal, M. D. and Miura, R. M.: Method for solving the Korteweg-de Vries equation, Phys. Rev. Lett. **19** 1095–1097 (1967)
- [8] Hatayama, G., Hikami, K., Inoue, R., Kuniba, A., Takagi T. and Tokihiro, T.: The $A_M^{(1)}$ automata related to crystals of symmetric tensors, *J. Math. Phys.* **42** 274–308 (2001)
- [9] Hatayama, G., Kuniba, A. and Takagi, T.: Soliton cellular automata associated with crystal bases, Nucl. Phys. B **577**[PM] 619–645 (2000)
- [10] Jimbo, M. and Miwa, T.: Solitons and infinite dimensional Lie algebras, Publ. RIMS. Kyoto Univ. **19** 943–1001 (1983)
- [11] Kac, M. and Moerbeke, P.: A complete solution of the periodic Toda problem, Proc. Nat. Acad. Sci. USA. **72** 2879–2880 (1975)
- [12] Kashiwara, M.: On crystal bases of the q -analogue of universal enveloping algebras. Duke Math. J. **63** 465–516 (1991)
- [13] Kirillov, A. N. and Reshetikhin, N. Yu.: The Bethe ansatz and the combinatorics of Young tableaux, J. Soviet Math. **41** 925–955 (1988)
- [14] Kuniba, A. and Nakanishi, T.: The Bethe equation at $q = 0$, the Möbius inversion formula, and weight multiplicities: I. The $sl(2)$ case, Prog. in Math. **191** 185–216 (2000)
- [15] Kuniba, A. and Sakamoto, R.: The Bethe ansatz in a periodic box-ball system and the ultradiscrete Riemann theta function, J. Stat. Mech. P09005 (2006)
- [16] Kuniba, A. , Sakamoto, R. and Yamada, Y.: Tau functions in combinatorial Bethe ansatz, preprint math.QA/0610505
- [17] Kuniba, A., Takagi, T. and Takenouchi, A.: Bethe ansatz and inverse scattering transform in a periodic box-ball system, Nucl. Phys. B [PM] 354–397 (2006)
- [18] Kuniba, A. and Takenouchi, A.: Bethe ansatz at $q = 0$ and periodic box-ball systems, J. Phys. A: Math. Gen. **39** 2551–2562 (2006)
- [19] Mumford, D.: Tata Lectures on Theta II, Birkhäuser, Boston (1984)
- [20] Takahashi, D. and Satsuma, J.: A soliton cellular automaton, J. Phys. Soc. Jpn. **59** 3514–3519 (1990)
- [21] Tokihiro, T., Takahashi, D., Matsukidaira and J. Satsuma, J.: From soliton equations to integrable cellular automata through a limiting procedure, Phys. Rev. Lett. **76** 3247–3250 (1996)
- [22] Yoshihara, D., Yura, F. and Tokihiro, T.: Fundamental cycle of a periodic box-ball system, J. Phys. A: Math. Gen. **36** 99–121 (2003)

Atsuo Kuniba:

Institute of Physics, Graduate School of Arts and Sciences, University of Tokyo,
Komaba, Tokyo 153-8902, Japan

`atsuo@gokutan.c.u-tokyo.ac.jp`

Reiho Sakamoto:

Department of Physics, Graduate School of Science, University of Tokyo, Hongo,
Tokyo 113-0033, Japan

`reiho@monet.phys.s.u-tokyo.ac.jp`